

Multicystic lesions of the spleen: A rare presentation of splenic lymphangioma in adult

Amarjothi J. M. V., Bennet D., Prabakaran R., Kannan D.,
Naganath Babu O. L.

ABSTRACT

Splenic lymphangiomas are rare tumors seen most commonly in children. However very rarely, they may also present as symptomatic multi cystic splenic lesions in adults also. Hence a high index of suspicion is needed to diagnose and treat these lesions when symptomatic and to prevent complications like rupture and infection. Accurate radiological diagnosis of splenic lymphangioma is difficult and histopathology with newer Immunohistochemistry markers like D240 is useful to diagnose splenic lymphangioma. We wish to present this interesting case of symptomatic splenic lymphangioma in a middle aged female patient.

Keywords: D240, Multi cystic spleen, Splenectomy, Splenic lymphangioma

Amarjothi J.M.V.¹, Bennet D.², Prabakar R.², Kannan D.³,
Naganath Babu O. L.⁴

Affiliations: ¹MS, MRCS, DNB, Registrar, Department of Surgical Gastroenterology, Madras Medical College, Chennai, India; ²MS, MCh, Assistant Professor, Department of Surgical Gastroenterology, Madras Medical College, Chennai, India; ³MS, MCh, Professor, Department of Surgical Gastroenterology, Madras Medical College, Chennai, India; ⁴MS, MCh, FRCS, MNAMS, Professor, Department of Surgical Gastroenterology, Madras Medical College, Chennai, India.

Corresponding Author: Amarjothi J. M. V., Room No 500, 4th floor, Tower 2, Department of Surgical Gastroenterology, MMC, Chennai-3, Tamil Nadu, India; Email: drmoses-vikramamarjothi@hotmail.com

Received: 12 April 2017
Accepted: 31 July 2017
Published: 28 June 2017

How to cite this article

Amarjothi JMV, Bennet D, Prabakar R, Kannan D, Naganath Babu OL. Multicystic lesions of the spleen: A rare presentation of splenic lymphangioma in adult. J Case Rep Images Pathol 2017;3:5–8.

Article ID: 100013Z11AV2017

doi: 10.5348/Z11-2017-13-CR-2

INTRODUCTION

Lymphangiomas, in general, are congenital malformations of the lymphatic system. Splenic involvement is rare in lymphangiomas with less than 1% of lymphangiomas localized intra-abdominally. Splenic lymphangiomas are usually benign and predominantly affect children, and only some cases have been reported in adults. Most cases of splenic lymphangioma in adults require surgical intervention as they may be symptomatic or lead to complications due to the natural course of the disease.

CASE REPORT

A 48-year-old female was admitted with multiple splenic cysts detected by abdominal ultrasonography, done for recurrent pain to the upper abdomen for two months. The pain was intermittent and localized to the epigastric and left hypochondrium. Physical examination detected moderate splenomegaly in the left upper quadrant extending for about 4 cm from the costal margin. Peripheral blood count, coagulation studies, and liver and kidney function tests were all within normal limits. Contrast-enhanced computed tomography scan showed

multiple non-enhancing cysts of variable size throughout the spleen (Figure 1A). Due to the presence of symptoms, large splenic size and probability of malignancy, open splenectomy was done. Spleen was found to be enlarged with multiple splenic cysts of variable size, where some contained clear fluid and others were hemorrhagic (Figure 1B). The histopathology was suggestive of splenic lymphangioma (Figure 2A) with multiple lymphatic vesicles of variable size between 1–3 cm, lined by a single layer of flat endothelial cells filled with clear mucinous fluid. Immunohistochemistry done was positive for Factor VIII related antigen (Factor VIII R Ag) (Figure 2B) implying the endothelial origin of the cells. D240, a novel IHC marker specific for lymphangioma was also found to be positive (Figure 2C).

DISCUSSION

Most common sites of occurrence of lymphangiomas include the neck, mediastinum and retroperitoneum [1]. Splenic lymphangiomas are benign cysts that commonly affect children, and are rarely reported in adults [2–4].




Figure 1: (A) Contrast-enhanced computed tomography showing numerous non-enhancing cystic lesions in an enlarged spleen (black arrows), (B) Splenectomy specimen showing numerous cysts in the spleen (white arrows).

In 1885, Frink reported the first lymphangioma in the spleen [5]. Most cases (80–90%) are reported in infants and children less than two years of age [6]. Cystic lesions of spleen include parasitic and non-parasitic cysts.

Among parasitic ones, echinococcal disease represents most of the cases [7]. Non-parasitic cysts are primary or true cysts and pseudocysts. Endothelial true cysts include lymphangiomas or hemangiomas [1]. Lymphangiomas can be classified as capillary, cavernous, or cystic. However, the delineation is not universally accepted [4, 8]. Cystic lymphangiomas are composed of numerous




Figure 2: (A) Multiple endothelium lined cystic spaces in splenic architecture (arrows) (H&E stain, x100), (B) Immunohistochemistry - Endothelial cells showing factor 8 related antigen positivity and granular reaction (arrows) (IHC, x200) (C) Immunohistochemistry D240 positivity conclusive of splenic lymphangioma (arrows).

cystic spaces containing amorphous eosinophilic fluid lined by a single layer of flat endothelial cells [7]. Splenic lymphangiomas are mostly asymptomatic and if symptomatic most commonly present with abdominal pain [8]. Diagnosis by radiology is usually not conclusive. In most cases, on CT scan, splenic lymphangiomas present with thin-walled cystic masses without enhancement or with only slight enhancement of the thin septa. Curvilinear peripheral mural calcifications are suggestive of cystic lymphangiomas but not specific [9]. On T1-weighted magnetic resonance imaging scan, the cystic lesions appear hypointense or hyperintense when filled with hemorrhagic or proteinaceous material [8, 9]. On T2-weighted images, they are multiloculated hyperintense corresponding to dilated lymphatic channels [8].

The differential diagnoses of splenic lymphangiomas include other solid and cystic lesions of the spleen, such as hemangioma, chronic infection, lymphoma and metastasis. Immunohistochemistry (IHC) markers including CD 31, CD 34, factor VIII R Ag, D240 and VEGFR-3 may show reactivity for splenic lymphangiomas. As CD34 for endothelial cells is not reliable, FVIII RAg and CD31 are used for identification of endothelial cells lining lymphatic spaces [5]. D240 is a monoclonal antibody initially developed against the M2A antigen or podoplanin, a fetal testis-related antigen [10]. Recently, D240 has been reported to be a selective marker for the lymphatic endothelium, and can be used for diagnosing splenic lymphangioma accurately from hemangioma also making it a valuable addition for studying benign and malignant vascular disorders in routinely processed tissue specimens [11].

VEGFR-3 is a marker for vascular endothelial growth factor C produced by the tumor cells which stimulate proliferation of adjacent lymphatics [12]. Electron microscopy, in splenic lymphangiomas, show rod-shaped storage granules of endothelial cells (Weibel Palade bodies), which emphasize the endothelial origin of these cells [13]. As these cysts are large and present with splenomegaly, they usually require splenectomy in most cases to prevent complications. The main complication is recurrence, which is demonstrated in 9.5% of patients, frequently after incomplete resection [5]. Other complications include rupture, infection, abscess formation, and pleural effusion or empyema, haemorrhage, consumptive coagulopathy, hypersplenism and even portal hypertension [14].

Partial resection techniques, aspiration, drainage and sclerosis are accompanied with high recurrence rates and are avoided. Laparoscopic splenectomy can be tried in patients with a normal size spleen.

CONCLUSION

Splenic lymphangioma should be considered in the differential diagnosis of a multicystic spleen irrespective of the age of diagnosis and may necessitate surgery including splenectomy for associated symptoms or to

prevent future complications. Histopathology and novel IHC markers like D240 are useful to make an accurate diagnosis of splenic lymphangiomas.

Author Contributions

Amarjothi J.M.V. – Substantial contributions to conception and design, Acquisition of data, Analysis and interpretation of data, Drafting the article, Revising it critically for important intellectual content, Final approval of the version to be published

Bennet D. – Substantial contributions to conception and design, Acquisition of data, Analysis and interpretation of data, Drafting the article, Revising it critically for important intellectual content, Final approval of the version to be published

Prabakar R. – Substantial contributions to conception and design, Acquisition of data, Analysis and interpretation of data, Drafting the article, Revising it critically for important intellectual content, Final approval of the version to be published

Kannan D. – Substantial contributions to conception and design, Acquisition of data, Analysis and interpretation of data, Drafting the article, Revising it critically for important intellectual content, Final approval of the version to be published

Naganath Babu O. L. – Substantial contributions to conception and design, Acquisition of data, Analysis and interpretation of data, Drafting the article, Revising it critically for important intellectual content, Final approval of the version to be published

Guarantor

The corresponding author is the guarantor of submission.

Conflict of Interest

Authors declare no conflict of interest.

Copyright

© 2017 Amarjothi J.M.V. et al. This article is distributed under the terms of Creative Commons Attribution License which permits unrestricted use, distribution and reproduction in any medium provided the original author(s) and original publisher are properly credited. Please see the copyright policy on the journal website for more information.

REFERENCES

1. Abbott RM, Levy AD, Aguilera NS, Gorospe L, Thompson WM. From the archives of the AFIP: primary vascular neoplasms of the spleen: Radiologic-pathologic correlation. *Radiographics* 2004 Jul-Aug;24(4):1137–63.
2. Vezzoli M, Ottini E, Montagna M, et al. Lymphangioma of the spleen in an elderly patient. *Haematologica* 2000 Mar;85(3):314–7.

3. Chang WC, Liou CH, Kao HW, Hsu CC, Chen CY, Yu CY. Solitary lymphangioma of the spleen: Dynamic MR findings with pathological correlation. *Br J Radiol* 2007 Jan;80(949):e4–6.
4. Beltrán MA, Barría C, Pujado B, et al. Gigantic cystic splenic lymphangioma. Report of one case. [Article in Spanish]. *Rev Med Chil* 2009 Dec;137(12):1597–601.
5. Solomou EG, Patriarheas GV, Mpadra FA, Karamouzis MV, Dimopoulos I. Asymptomatic adult cystic lymphangioma of the spleen: Case report and review of the literature. *Magn Reson Imaging* 2003 Jan;21(1):81–4.
6. Bardeguez A, Chatterjee M, Tepedino M, Sicuranza B. Systemic cystic angiomas in pregnancy: A case presentation and review of the literature. *Am J Obstet Gynecol* 1990 Jul;163(1 Pt 1):42–5.
7. Lee WS, Kim YH, Chee HK, Lee SA, Kim JD, Kim DC. Cavernous lymphangioma arising in the chest wall 19 years after excision of a cystic hygroma. *Korean J Thorac Cardiovasc Surg* 2011 Oct;44(5):380–2.
8. Ogun GO, Oyetunde O, Akang EE. Cavernous lymphangioma of the breast. *World J Surg Oncol* 2007 Jun 20;5:69.
9. Urrutia M, Mergo PJ, Ros LH, Torres GM, Ros PR. Cystic masses of the spleen: Radiologic-pathologic correlation. *Radiographics* 1996 Jan;16(1):107–29.
10. Schacht V, Dadras SS, Johnson LA, Jackson DG, Hong YK, Detmar M. Up-regulation of the lymphatic marker podoplanin, a mucin-type transmembrane glycoprotein, in human squamous cell carcinomas and germ cell tumors. *Am J Pathol* 2005 Mar;166(3):913–21.
11. Kahn HJ, Bailey D, Marks A. Monoclonal antibody D2-40, a new marker of lymphatic endothelium, reacts with Kaposi's sarcoma and a subset of angiosarcomas. *Mod Pathol* 2002 Apr;15(4):434–40.
12. Debelenko LV, Perez-Atayde AR, Mulliken JB, Liang MG, Archibald TH, Kozakewich HP. D2-40 immunohistochemical analysis of pediatric vascular tumors reveals positivity in kaposiform hemangioendothelioma. *Mod Pathol* 2005 Nov;18(11):1454–60.
13. Takayama A, Nakashima O, Kobayashi K, Kojiro M. Splenic lymphangioma with papillary endothelial proliferation: A case report and review of the literature. *Pathol Int* 2003 Jul;53(7):483–8.
14. Morgenstern L, Bello JM, Fisher BL, Verham RP. The clinical spectrum of lymphangiomas and lymphangiomatosis of the spleen. *Am Surg* 1992 Oct;58(10):599–604.

Access full text article on
other devices



Access PDF of article on
other devices

